

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Offic

An agency of
Industry Canada

= EP 0514 769
BCS 02-1006
CA 2068826 C 2003/01/07
BI (11)(21) 2 068 826

(12) BREVET CANADIEN
CANADIAN PATENT

(13) C

(22) Date de dépôt/Filing Date: 1992/05/15

(41) Mise à la disp. pub./Open to Public Insp.: 1992/11/19

(45) Date de délivrance/Issue Date: 2003/01/07

(30) Priorité/Priority: 1991/05/18 (P 41 16 440.7) DE

(51) Cl.Int.⁵/Int.Cl.⁵ A01N 47/36, A01N 47/30, A01N 25/30,
A01N 25/04

(72) Inventeurs/Inventors:
FRISCH, GERHARD, DE;
MAIER, THOMAS, DE

(73) Propriétaire/Owner:
HOECHST AKTIENGESELLSCHAFT, DE

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : SUSPENSION-EMULSION A BASE DE FENOXAPROP-ETHYLE NOUVEAU

(54) Title: NOVEL SUSPOEMULSIONS BASED ON FENOXAPROP-ETHYL

(57) Abrégé/Abstract:

The invention relates to storage-stable aqueous suspoemulsions which comprise fenoxaprop-ethyl, a sulfonylurea and/or phenylurea, an aromatic solvent and a surfactant combination of ethoxylated tristyrylphenol and ethoxylated synthesis alcohol, which are phosphated if appropriate, to processes for their preparation and to their use in plant protection.

Canada

<http://opic.gc.ca> • Ottawa-Hull K1A 0C9 • <http://cipo.gc.ca>

OPIC • CIPQ 191

OPIC



CIPQ

Description

Novel suspoemulsions based on fenoxaprop-ethyl

The invention relates to suspoemulsions based on fenoxaprop-ethyl. Suspoemulsions (SE) of various types are described, for example, in EP-A-0117999, U.S. Patent 4824663, EP-A-0289356, EP-A-0261492 and EP-A-0143099. A review of the latest art in the field of suspoemulsions is to be found on the one hand in Pestic. Sci. 1990, 29, 451 - 465 (Recent Development in Suspoemulsions, P. Mulqueen et al.) and on the other hand in Pestic. Sci. 1990, 29, 437 - 449 (Trends in the Formulation of Pesticides - An Overview, D. Seaman). It is furthermore known that a formulation depends very much on the active compounds contained therein. The influence of the active compounds then manifests itself quite decisively in the nature and composition of surfactants in these multiphase formulations. The corresponding solvent for the phase which is to be emulsified and in which one or more active compounds are dissolved also plays an important role here.

Already existing, known systems - such as, for example, those mentioned above - therefore cannot always be resorted to for the preparation of suspoemulsions of novel active compounds, such as, for example, fenoxaprop-ethyl, with representatives from the novel class of sulfonylureas, such as, for example, amidosulfuron, or the already known phenylurea derivatives, such as, for example, isoproturon. Fenoxaprop-ethyl thus does not have too good a solubility in phthalic acid esters (EP-A-0177999), while it has a very good solubility in aromatic solvents such as xylene or ethylbenzene, or in mixtures of aromatics, such as, for example, the solvents of the "Solvesso" series from Esso. Nevertheless, no particularly stable emulsion can be formed from fenoxaprop-ethyl in such solvents with block polymers (EP-A-0261492) or "graft copolymers" (EP-A-0289356 or GB Patent 2026341 A) in dispersion-containing water.

It has now been found, surprisingly, that storage-stable and technologically problem-free suspoemulsions of fenoxaprop-ethyl with herbicides from the sulfonylurea series, for example amidosulfuron, or from the phenylurea series, for example isoproturon, can be prepared with the aid of a certain solvent and surfactant combination. Moreover, in the case of amidosulfuron virtually no chemical degradation of the active compound is found, although this could be expected on the basis of the chemical structure. Phenylurea derivatives, such as isoproturon, are stable in aqueous dispersions. Safeners can also be added to these suspoemulsions. This also particularly applies to fenoxaprop-ethyl if, for example, it is dissolved in the organic phase in combination with the safener ethyl 1-(2,4-dichlorophenyl)-5-trichloromethyl-(1H)-1,2,4-triazole-3-carboxylate (EP-A-0174562). No interactions which have adverse effects on the stability and action of the system take place between the safener and the fenoxaprop-ethyl in the system described.

The invention therefore relates to suspoemulsions which comprise fenoxaprop-ethyl, at least one herbicidal active compound from the sulfonylurea series and/or from the phenylurea series, an aromatic solvent or solvent mixture and a surfactant combination of ethoxylated tristyrylphenol and ethoxylated sterically modified synthesis alcohol of average chain length C_{13} , each of which can be phosphated and neutralized with alkali or amine.

Fenoxaprop-ethyl can be employed as a stereoisomer mixture (for example as the racemate) or in the form of the D(+)-isomer fenoxaprop-P-ethyl.

Possible herbicides from the sulfonylurea series are pyrimidine- or triazinylaminocarbonyl-[benzene-, pyridine-, pyrazole-, thiophene- and (alkylsulfonyl)-alkylamino-]-sulfamides. Preferred substituents on the pyrimidine ring or triazine ring are alkoxy, alkyl,


- haloalkoxy, haloalkyl, halogen or dimethylamino, it being possible for all the substituents to be combined independently of one another. Preferred substituents in the benzene, pyridine, pyrazole, thiophene or (alkylsulfonyl)alkylamino part are alkyl, alkoxy, halogen, nitro, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkoxyaminocarbonyl, alkyl, alkoxyaminocarbonyl, haloalkoxy, haloalkyl, alkylcarbonyl, alkoxyalkyl and (alkanesulfonyl)alkylamino.
- 10 Examples of suitable sulfonylureas are
- 1) phenyl- and benzylsulfonylureas and related compounds, for example 1-(2-chlorophenylsulfonyl)-3-(4-methoxy-6-methyl-1,3,5-triazin-2-yl)-urea (chlorsulfuron), 1-(2-ethoxycarbonylphenylsulfonyl)-3-(4-chloro-6-methoxy-15 pyrimidin-2-yl)-urea (chlorimuron-ethyl), 1-(2-methoxyphenylsulfonyl)-3-(4-methoxy-6-methyl-1,3,5-triazin-2-yl)-urea (metsulfuron-methyl), 1-(2-chloroethoxy-phenylsulfonyl)-3-(4-methoxy-6-methyl-1,3,5-triazin-2-yl)-urea (triasulfuron), 20 1-(2-methoxycarbonyl-phenylsulfonyl)-3-(4,6-dimethylpyrimidin-2-yl)-urea (sulfometuron-methyl), 1-(2-methoxycarbonylphenylsulfonyl)-3-(4-methoxy-6-methyl-1,3,5-triazin-2-yl)-3-methylurea (tribenuron-methyl)
 - 25 1-(2-methoxycarbonylbenzylsulfonyl)-3-(4,6-dimethoxy-pyrimidin-2-yl)-urea (bensulfuron-methyl) 1-(2-methoxycarbonylphenylsulfonyl)-3-(4,6-bis-(difluoromethoxy)-pyrimidin-2-yl)-urea (primisulfuron-methyl), 3-(4-ethyl-6-methoxy-1,3,5-triazin-2-yl)-1-(2,3-dihydro-30 1,1-dioxo-2-methylbenzo[b]thiophene-7-sulfonyl)-urea (see EP-A-79683) and 3-(4-ethoxy-6-ethyl-1,3,5-triazin-2-yl)-1-(2,3-dihydro-1,1-dioxo-2-methylbenzo[b]thiophene-7-sulfonyl)-urea (see EP-A-79683),

28976-46

-4-

- 2) thienylsulfonylureas, for example
 1-(2-methoxycarbonylthiophen-3-yl)-3-(4-methoxy-6-methyl-1,3,5-triazin-2-yl)-urea (thifensulfuron-methyl),
- 3) pyrazolylsulfonylureas, for example
 1-(4-ethoxycarbonyl-1-methylpyrazol-5-yl-sulfonyl)-3-(4,6-dimethoxypyrimidin-2-yl)-urea (pyrazosulfuron-methyl) and methyl 3-chloro-5-(4,6-dimethoxypyrimidin-2-ylcarbamoylsulfamoyl)-1-methyl-pyrazole-4-carboxylate (see EP-A-282613),
- 4) sulfonyldiamide derivatives, for example
 3-(4,6-dimethoxypyrimidin-2-yl)-1-N-methyl-N-methylsulfonylamidosulfonyl)-urea (amidosulfuron) and structural analogs (see EP-A-0131258 and Z. Pfl. Krankh. Pfl. Schutz, Special Edition XII, 489-497 (1990)),
- 5) pyridylsulfonylureas, for example 1-(3-N,N-dimethylaminocarbonylpyridin-2-yl-sulfonyl)-3-(4,6-dimethoxypyrimidin-2-yl)-urea (nicosulfuron), 1-(3-ethylsulfonylpyridin-2-yl-sulfonyl)-3-(4,6-dimethoxypyrimidin-2-yl)-urea (DPX-E 9636, see Brighton Crop Prot. Conf. - Weeds - 1989, page 23 et seq.) and pyridylsulfonylureas such as are described in German Patent Applications Publication Nos. DE-A 4000503 and DE-A 4030577, preferably those of the formula I or salts thereof,

25




(I)

in which

E is CH or N, preferably CH,

- R^8 is iodine or $NR^{13}R^{14}$,
 R^9 is hydrogen, halogen, cyano, (C_1-C_3) -alkyl, (C_1-C_3) -alkoxy, (C_1-C_3) -haloalkyl, (C_1-C_3) -haloalkoxy, (C_1-C_3) -alkylmercapto, (C_1-C_3) -alkoxy- (C_1-C_3) -alkyl, (C_1-C_3) -alkoxy-carbonyl, mono- or di- (C_1-C_3) -alkyl-amino, (C_1-C_3) -alkyl-sulfinyl or -sulfonyl, $SO_2-NR^aR^b$ or $CO-NR^aR^b$, in particular H,
 R^a and R^b independently of one another are hydrogen, (C_1-C_3) -alkyl, (C_1-C_3) -alkenyl, (C_1-C_3) -alkynyl or together $-(CH_2)_4-$, $-(CH_2)_5-$ or $(CH_2)_2-O-(CH_2)_2-$,
 R^{10} is H or CH_3 ,
 R^{11} is halogen, (C_1-C_2) -alkyl, (C_1-C_2) -alkoxy, (C_1-C_2) -haloalkyl, preferably CF_3 , or (C_1-C_2) -haloalkoxy, preferably $OCHF_2$ or OCH_2CF_3 ,
 R^{12} is (C_1-C_2) -alkyl, (C_1-C_2) -haloalkoxy, preferably $OCHF_2$, or (C_1-C_2) -alkoxy and
 R^{13} is (C_1-C_4) -alkyl and R^{14} is (C_1-C_4) -alkylsulfonyl, or R^{13} and R^{14} together are a chain of the formula $-(CH_2)_3SO_2-$ or $-(CH_2)_4SO_2-$, for example 3-(4,6-dimethoxy-2-pyrimidin-2-yl)-1-(3-N-methylsulfonyl-N-methylaminopyridin-2-yl)-sulfonylurea, and

6) alkoxyphenoxy-sulfonylureas such as are described in EP-A-0342569, preferably those of the formula II or salts thereof,



- in which
 E is CH or N, preferably CH,
 R^{15} is ethoxy, propoxy or isopropoxy,
 R^{16} is hydrogen, halogen, nitro, CF_3 , CN, (C_1-C_4) -alkyl, (C_1-C_4) -alkoxy, (C_1-C_4) -alkylmercapto or

- (C₁-C₃)-alkoxy-carbonyl, preferably in the 6-position on the phenyl ring,
n is 1, 2 or 3, preferably 1,
R¹⁷ is hydrogen, (C₁-C₄)-alkyl or (C₃-C₄)-alkenyl,
5 R¹⁸ and R¹⁹ independently of one another are halogen, (C₁-C₂)-alkyl, (C₁-C₂)-alkoxy, (C₁-C₂)-haloalkyl, (C₁-C₂)-haloalkoxy or (C₁-C₂)-alkoxy-(C₁-C₂)-alkyl, preferably OCH₃ or CH₃, for example 3-(4,6-dimethoxypyrimidin-2-yl)-1-(2-ethoxy-
10 phenoxy)-sulfonylurea,
and other related sulfonylurea derivatives and mixtures thereof. Amidosulfuron and/or isoproturon are preferred.

- Examples of suitable phenylurea derivatives are isoproturon, diuron, chlortoluron, monolinuron, linuron, neburon,
15 monuron, fluometuron, fenuron, siduron, terbuthuron, chlorbromuron and tetrafluoron.

The content of fenoxaprop-ethyl is preferably 0.1 to 20 % by weight, in particular 0.2 to 10 % by weight.

- The content of sulfonylurea derivatives and/or phenylurea
20 derivatives, preferably of sulfonyldiamide derivatives or isoproturon, is in general preferably 0.1 to 60 % by weight, in particular 1 to 45 % by weight, amidosulfuron in particular also being employed in somewhat smaller amounts (preferably 0.1 to 30 % by weight, in particular
25 0.2 to 15 % by weight).

- Possible preferred solvent combinations are aromatic solvents which are derived from benzene, such as xylene, mesitylene, indane, diisopropylbenzene and higher homologs, as well as solvents from the "Solvesso series from
30 Esso or a mixture of the solvents mentioned. A naturally occurring oil of animal or vegetable origin is preferably added to these solvents or solvent mixtures, it being possible for the ratio of oil to aromatic to be 1:1 to 1:100, but preferably 1:4 to 1:50. The addition of

naturally occurring oils not only brings about an improvement in the shear stability of the system, which is essential for preparation and storage, but can also additionally increase the penetration properties. The
5 addition of a vegetable oil, such as rapeseed oil, is preferred. The total solvent content is preferably 2 to 70 % by weight, in particular 3 to 50 % by weight.

A possible surfactant mixture according to the invention is a combination of ethoxylated tristyrylphenols and
10 ethoxylated, sterically modified synthesis alcohol of average chain length C_{13} (isotridecyl alcohol), which can also be phosphated and neutralized with alkali or amines. 0.1 to 30 % by weight, preferably 0.5 to 20 % by weight, of the surfactant mixture is added. The degree of ethox-
15 ylation in the case of the tristyrylphenol derivatives can be between 10 and 40, but preferably between 14 and 28. The degree of ethoxylation of the isotridecyl alcohol can be between 2 and 20, preferably between 4 and 14. The tristyrylphenol derivatives are marketed, for
20 example, under the trade names °Soprophor BSU, °Soprophor 3D33, °Soprophor FL, °Soprophor CY/8, °Soprophor S/25, °Soprophor FL/60, °Soprophor FLK (Rhone Poulenc), Hoe S 3474, Hoe S 3475 and Hoe S 3775; the isotridecyl alcohol derivatives are from the °Genapol X
25 series (Hoechst). The ratio of ethoxylated tristyrylphenol derivatives to ethoxylated isotridecyl alcohol derivatives is 25:1 to 1:35, preferably 4:1 to 1:8.

Formulations which contain only tristyrylphenol deriva-
tives, that is to say not the surfactant mixture accord-
30 ing to the invention, are not stable to shear in the present case and have a marked tendency to agglomerate (see examples).

Another unexpected advantage is that this surfactant combination is accompanied by a significant improvement
35 in the biological action. Experiments have shown

(Table II) that, for example in the case of the combination isoproturon:fenoxaprop-ethyl, formulations with the above surfactant mixture were significantly better than comparable tank mixtures with the commercially available individual formulations or if, for example, ethoxylated isotridecyl alcohol was replaced by similar wetting agents, such as °Genapol LRO.

The formulations according to the invention can comprise a dodecyl- or tridecylbenzenesulfonate in an amount of 0.01 - 12 % by weight as an additional surfactant which helps to improve the properties of dispersal of the stably suspended particle but is not essential. Thus, for example, °Maranil (dodecylbenzenesulfonate Na salt) from Henkel can be employed as a paste or powder.

Possible additional surfactants are furthermore those polyacrylic acid derivatives such as can be obtained, for example, under the trade names °Sokalan CP10 (BASF), the °Geropon series (HB, DA, DG) (Rhone Poulenc) or the °Dispersant series (Rhone Poulenc) or the °Degapas series (Degussa).

°Sokalan CP10 is a modified Na polyacrylate of low molecular weight, which is prepared by a special polymerization process (BASF Techn. Info TI/P 3039 d of 1988.).

The °Geropon types HB, DA and DG and °Dispersant HB and FB are, according to the Rhone-Poulenc data sheet of 1979 and 1989, alkali metal polyacrylates which are available both in liquid and in solid form.

The °Degapas series are also alkali metal or ammonium salts of polyacrylic acid derivatives.

Up to 25 % by weight, preferably up to 15 % by weight, of commercially available auxiliaries, such as wetting agents, dispersing agents, foam suppressants, thickeners,

preservatives and antifreeze agents, can furthermore also additionally be added.

5 Examples of possible additional wetting and dispersing agents are tributylphenol polyglycol ethers, such as the °Sapogenat T brands (Hoechst) or nonylphenol polyglycol ethers, such as the °Arkopal N brands (Hoechst).

Suitable foam suppressants are, for example, those based on silicone, such as those from the °Silcolapse series (Rhone Poulenc), SE 39 or antifoam agent SH (Wacker).

10 Thickeners can be inorganic or organic in nature; they can also be combined. Suitable thickeners are, for example, those based on aluminum, xanthan, methylcellulose, polysaccharide, alkaline earth metal silicate, gelatine and polyvinyl alcohol, such as, for example,
15 °Bentone EW, °Veegum, °Rhodopol 23 or °Kelzan S.

If necessary, preservatives are used, for example those based on formaldehyde, benzoic acid and triphenyltin, such as, for example, °Kobate C.

20 Antifreeze agents, such as urea, salts, polyols (for example glycol, propylene glycol or glycerol) or sugars, can furthermore also be added.

The invention furthermore relates to a method of combating undesirable plant growth, which comprises applying a formulation according to the invention to plants, plant
25 seeds or the cultivation area.

Suspoemulsions according to the invention are advantageously prepared from the individual dispersions of the dispersed active compounds, such as are used, for example, in the case of the phenylurea derivatives, such
30 as, for example, linuron, isoproturon, diuron and chlorotoluron and mixtures thereof. The active compounds are

28976-46

-10-

employed as finished formulations with the customary formulation auxiliaries, such as are described, for example in EP-A-0022925, EP-A-0110174 and German Patent Application Publication No. DE-A- 3538247. The suspoemulsion can be
5 mixed intimately by, for example, wet grinding.

Aqueous dispersions are similarly also used as starting substances in the case of the sulfonylurea derivatives, such as, for example, amidosulfuron, as has been proposed in German Patent Application Publication No.
10 DE-A-411644.5 (EP-A-0514768). The aqueous dispersion phase can be brought together with the organic active compound solution phase, for example, in accordance with EP-A-0130370.

The weight ratio of aqueous to organic phase can
15 be 1:1.5 to 50:1, but preferably 1:1 to 35:1.

Examples of the novel suspoemulsions according to the invention are listed in the following Tables I a and I b, without the invention being limited thereto.

[illegible]

T A B L E I b

Example	14	15	16	17	18	19	20	21	22	23
Amidosulfuron	3.26	3.26	3.23	3.4	3.6	3.2	3.2	3.2	3.5	3.5
Fenoxaprop-P-ethyl	7.13	7.13	7.06	7.3	6.92	6.92	7.0	7.0	8.0	8.0
Safener ¹⁾					3.06	3.06	3.1	3.1	3.7	3.7
*Solvesso 150	32.87	32.87	32.94	33.43	22.02	22.0	22.0	22.0	25.5	25.5
Rapeseed oil	5.0	5.0	5.0	5.0	4.0	4.0	4.0	4.0	4.0	4.0
*Soprophor BSU	5.0	5.0			4.0			4.0		
*Soprophor FL			5.0			4.0	4.0			4.0
Hoe S 3474				5.0					4.5	
*Sokalan CP10	1.27	1.27	1.27	1.3	1.3	1.3	1.3	1.3	1.5	1.7
Glycerol	1.0	2.0	1.0	1.02	1.0	1.0	2.0	1.0	2.0	1.0
*Maranil A	0.25	0.5	0.5	0.5	0.5	0.25	0.5	0.25	0.25	0.5
*Darvan No. 3	0.127	0.127	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13
SE 39	0.127	0.127	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13
*Silcolapse 5020	0.127	0.127	0.127	0.127	0.127	0.13	0.13	0.13	0.13	0.13 (cont.)

*Rhodopol 23	0.025	0.025	0.025	0.025	0.025	0.025	0.025	0.025	0.025
*Kobate C	0.013	0.013	0.013	0.013	0.013	0.015	0.013	0.013	0.013
Water to 100 %									

Ethyl 1-(2,4-dichlorophenyl)-5-trichloromethyl-(1H)-1,2,4-triazole-3-carboxylate
Darvan No. 3 is the sodium salt of a polymeric benzoylalkylsulfonic acid

T A B L E I I
Biological comparison experiments

	Herbicidal action against wild oats	
Tank mix of fenoxaprop- P-ethyl + isoproturon 500 g/l	(0.5 + 1 l/ha)	standard
Example 7	(1.25 l/ha)	+ (better than standard)
Example 9	(1.25 l/ha)	++ (significantly better than standard)

28976-46

-15-

CLAIMS:

1. A suspoemulsion comprising 0.1 to 20% by weight of fenoxaprop-ethyl as a stereoisomer mixture or as the D(+)-isomer, 0.1 to 60% by weight of at least one herbicidal
5 active compound from the sulfonylurea series or the phenyl urea series, 2 to 70% by weight of an aromatic solvent or solvent mixture, and 0.1 to 30% by weight of a surfactant combination of ethoxylated tristyrylphenol and ethoxylated sterically modified synthesis alcohol of average chain
10 length C₁₃, each of which can be phosphated and neutralized with alkali or amine, wherein the ratio of ethoxylated tristyrylphenol and ethoxylated sterically modified synthesis alcohol is 25:1 to 1:35.
2. A suspoemulsion as claimed in claim 1 comprising
15 0.2 to 10% by weight of fenoxaprop-ethyl as a stereoisomer mixture or as the D(+)-isomer, 1 to 45% by weight of at least one herbicidal active compound from the sulfonylurea series or the phenyl urea series, 3 to 50% by weight of an aromatic solvent or solvent mixture, and 0.5 to 20% by
20 weight of the surfactant combination, wherein the ratio of ethoxylated tristyrylphenol and ethoxylated sterically modified synthesis alcohol is 4:1 to 1:8.
3. A suspoemulsion as claimed in claim 1 or 2, which comprises a solvent mixture of an aromatic solvent and a
25 naturally occurring oil of animal or vegetable origin.
4. A suspoemulsion as claimed in any one of claims 1 to 3, which additionally comprises another surfactant and/or customary auxiliaries from the series comprising wetting agents, dispersing agents, foam suppressants, thickeners,
30 preservatives and antifreeze agents.

28976-46

-16-

5. A suspoemulsion as claimed in any one of claims 1 to 4, comprising amidosulfuron and/or isoproturon.

6. A process for the preparation of a suspoemulsion as claimed in any one of claims 1 to 5, which comprises
5 bringing together dispersions of the individual active compounds and mixing or homogenizing them mechanically with the other constituents of the suspoemulsion.

7. The use of a suspoemulsion as claimed in any one of claims 1 to 5 for combating undesirable plant growth.

10 8. A method of combating undesirable plant growth, which comprises applying a suspoemulsion as claimed in any one of claims 1 to 5 to plants, plant seeds or the cultivation area.

9. A composition consisting essentially of 0.1 to 20%
15 by weight of fenoxaprop-ethyl as a stereoisomer mixture or as the D(+)-isomer, 0.1 to 60% by weight of at least one herbicidal active compound from the sulfonylurea series or the phenylurea series, 2 to 70% by weight of an aromatic solvent or solvent mixture, and 0.1 to 30% by weight of a
20 surfactant combination of ethoxylated tristyrylphenol and ethoxylated sterically modified synthesis alcohol of average chain length C₁₃, each of which can be phosphated and neutralized with alkali or amine, wherein the ratio of ethoxylated tristyrylphenol and ethoxylated sterically
25 modified synthesis alcohol is 25:1 to 1:35; and, the composition is a suspoemulsion.

FETHERSTONHAUGH & CO.
OTTAWA, CANADA

PATENT AGENTS

Abstract of the disclosure:

Novel suspoemulsions based on fenoxaprop-ethyl

The invention relates to storage-stable aqueous suspoemulsions which comprise fenoxaprop-ethyl, a sulfonylurea and/or phenylurea, an aromatic solvent and a surfactant combination of ethoxylated tristyrylphenol and ethoxylated synthesis alcohol, which are phosphated if appropriate, to processes for their preparation and to their use in plant protection.